

## California Organic **FERTILIZERS**

## PHYTA-CAL QC **Liquid Calcium 8%**

## **Guaranteed Analysis**

Calcium (Ca)...... 8.0%

Derived from Calcium Chloride.

### CAUTION: Keep Out of Reach of Children

#### WARNING

Harmful if swallowed. Avoid breathing spray mist.

May cause irritation of nose, throat, and/or skin. Avoid contact with eyes, skin, and clothing. If swallowed: Induce vomiting by sticking finger down throat. Never give anything by mouth to an unconscious person. Get medical attention If in eyes: Wash eyes with plenty of clear water to at least 15 minutes. Get medical attention.

> Information regarding the contents and levels of metals in this product is available on the Internet at https://www.aapfco.org/metals.htm

CONDITIONS OF SALE: \( \). Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when CONDITIONS OF SALE: used in accordance with directions under normal conditions of use. No one other than an officer of seller, is authorized to make any warranty, guarantee, or direction concerning this product. 2. Because the time, place, and rate of application are beyond seller's control, seller's liability from handling, storage, and use of this product is limited to replacement of product or refund of purchase price.

## CALIFORNIA ORGANIC FERTILIZERS, INC.

10585 Industry Ave. Hanford, California 93230 (800) 269-5690 · Fax: (559) 582-2011 www.organicag.com Rev. 5-19-22

Lot #: \_\_\_\_\_



#### **GENERAL INFORMATION**

PHYTA-CAL QC<sup>™</sup> can be tank mixed and applied with most commonly used insecticides and fungicides. DO NOT tank mix with growth hormone products, plant growth regulators, or foliar fertilizers unless specifically recommended

#### DIRECTIONS FOR USE

Mix well before use. It is more effective to spray small amounts with greater frequency. DO NOTUSE
PHYTA-CAL QC™ WITH OTHER FOLIAR PERTILIZERS
AS PRECIPITATION MAY OCCUR

Cotton, Soybeans, Beans, Tomatoes, Melons, Cucurbits, Peanuts, Peppers

Begin application when first flower appears. Spray enough water to achieve thorough coverage of the entire plant. Spray schedule may be discontinued when when last fruit, boll, or pods are set. Use 2 to 4 quarts per treated acre at 7 to 10 day intervals.

# Celery, Carrots, Beets, Cole Crops,

Potatoes, Sweet Potatoes
Begin application when the crop begins to grow rapidly. This will be after thinning on most row crops. Spray enough water to achieve thorough coverage of the entire plant. Spray schedule may be discontinued when rapid growth ceases. Use 2 to 4 quarts per treated acre at 7 to 10 day intervals.

#### Lettuce, Leaf Crops

Begin treatment after thinning and continue up to one week before harvest. Use 1 to 2 quarts per acre per application. Treat at 7 to 10 day intervals.

#### Citrus Fruits, Nuts, Apples, Peaches, Pears, Cherries, Almonds

Begin application when buds first form. Spray enough water to achieve thorough coverage of the entire tree. Spray must be discontinued when fruit has sized. Use 2 to 4 quarts per acre. Low volume application is preferred.

#### Grapes

Begin application when fruit first starts to form. Spray enough water to achieve thorough coverage of the entire plant canopy. Spray must be discontinued when grapes are sized. Use 2 to 4 quarts per acre at 7 to 10 day intervals. A 4 quart dormant application is also recommended

PHYTA-CAL QC<sup>™</sup> is recommended on the following crops: alfalfa, almonds, apples, avocados, barley, all beans, broccoli, cabbage, carrots, cauliflower, celery, citrus, corn, grapes, lettuce, melons, milo, nectarines, peaches, peanuts, pears, pecans, peppers, potatoes, rice, sorghum, soybeans, strawberries, sugar beets, sugar cane, sweet corn, tomatoes, turnips, walnuts, watermelons, wheat, and most other crops.

Package Size:	☐ 275 gal. / 1041 L
☐ 5 gal. / 18.9 L ☐ 55 gal. / 208.2 L	☐ Bulk ☐ Sample

Density: 10.5 lbs. per gallon at 68° F